

PEANUT'S MEDICAL HISTORY

A5757645 66.60LBS DOG TAN N PIT BULL

2/24/2026

INTAKE EXAM

NORMAL

66.60LBS

Treatment Description:

Dog FAS Intake Medical/Behavior Form

S:

Patient was brought in as Owner surrender , OTC

Microchip not found

O:

Behavioral Observations

Offer tasty food: No interest

Type of food offered: Canned

Overall body posture: Upright , Stiff/tense

Ears: Neutral

Eyes: Hard stare

Mouth: Lips tense/short , Lip lift/showing teeth

Vocalizing: Growling

Fear-Anxiety-Stress level observed at intake: (5) High FAS

Other behavior observations: was originally removed from kennel with a leash but then began lunging and trying to bite, come along and vaccine gate needed, started alligator rolling on come along

Medical Observations:

WT 66.6 lbs

Age: 1 year

Gender: male

BCS 4 / 9

MM- pink CRT- uto BAR

EENT: unable to examine

CV: uto

RESP: uto

Abdomen/UG: visable testicles

PLN: uto

Musculoskeletal: amb x 4

Integument: NSF

Neuro: uto

P: Routine vaccinations:

Nobivac DHLPP SQ RFL (booster in 2-4 weeks)

Bordetella Intratrach3 IN once (good for 1 year)

Bravecto 44-88lbs PO once

6.7 cc Pyrantel pamoate suspension 50mg/ml PO once

Examined b

Assisted by:

6.70 DEWORM PYRANTEL 1.00
TIMES/DAY FOR 1.00 DAYS

Date Given:2/24/2026

BRAVECTO CANINE 44-88LBS

Date Given:2/24/2026

BORDETELLA

Date Given:2/24/2026

DHLPP

Date Given:2/24/2026

3/19/2026

SX EXAM

NORMAL

55.40LBS

Treatment Description:

NOBIVAC 3 Serial number: 838919 eXP 10/31/26

Microchip applied # 956000019945264

Rx Vetprofen 100 mg tablet SID for 3 days (Pain Medication)

S: Patient presents for Dog Castration.

O: Physical examination

Temperature- 101.2 Pulse- 120 Respiration- 50 MM- pink CRT- < 2 sec BAR

Weight 55.4 lbs BCS 5/9

EENT- mild dental calc.

Cardiovascular/Respiratory- NSF

Abdomen- NSF

Musculoskeletal- NSF

Neurologic- NSF

Lymph nodes- NSF

Genito-urinary- NSF

Integument- thinning coat on dorsal lumbar, irritation on right side of scrotal sac

Behavior: tense, nervous, stiff, no aggression, used muzzle out of precaution.

A: Patient is fit for surgery

P: Surgery Report- Dog Neuter

Anesthesia

Pre-anesthetic: Acepromazine (2mg/ml) 0.6 cc IM, Butorphanol 10mg/ml: 0.75 cc IM,

Ampicillin 200mg/ml: 2.8 cc SQ

Induction: Ketamine (100 mg/ml) 1.2 cc IV, Midazolam (5 mg/ml) 1.2 cc IV

Intubation: Endotracheal tube size 10

Maintenance: Oxygen 1L Isoflurane 2%

Monitoring: Pulse oximeter

Supportive care: Thermoregulation pad/blanket

A testicle was pushed up in the scrotum on the median raphe area and a slightly off midline incision was made over the testicle (#10). The testicle was then exteriorized and freed of subcutaneous and connective tissue, manually with gauze. A Carmalt was then placed over the vas deferens, cremaster, and vessels proximal to the epididymus. A single ligature (Miller's knot, 2-0 PDS) was placed proximal to the clamp and the testicle removed distal to the clamp. The clamp was then removed and the stump checked for bleeding. This procedure was repeated on the second testicle. The subcutaneous tissue was apposed using a single cruciate suture of 2-0 Monocryl. Tissue glue was placed on the skin.

Intradermal tattoo placed

Surgeon

VETPROFEN 100

Date Given:3/19/2026

RABIES

Date Given:3/19/2026

A5757645 55.40LBS DOG TAN N PIT BULL

IF YOU OR YOUR VETERINARIAN HAVE ANY QUESTIONS REGARDING THE MEDICAL TREATMENT YOUR
ANIMAL HAS RECEIVED
PLEASE CALL US AT (562) 940-6898